

Clinician-rated Level 1 Cross-cutting Measures

Clinician-rated Level 1 Cross-cutting Measures of Psychosis and Suicide Risk

Level 1 Clinician Rating Scale for Psychosis					
Presence of Psychosis:	Not present	Equivocal	Present, but mild	Present and moderate	Present and severe
Psychosis: As manifested over the last 2 weeks, by delusions, hallucinations, or disorganized speech	0	1	2	3	4

Level 1 Clinician Rating of Current Concern about Potential Suicide Behavior

Level 1 Clinician Rating of Current Concern About Potential Suicide Behavior:

Note: The items below should serve as a guideline of minimum factors to be considered when determining your concern about the patient's potential risk for suicidal behaviors.

Is there evidence of:

LONG-TERM FACTORS:

1. Any history of suicide attempt?

- No
 Yes

2. Any history of mental illness?

- No
 Yes

3. Any history of physical or sexual abuse?

- No
 Yes

4. Long-standing tendency to lose temper or become aggressive with little provocation?

- No
 Yes

5. Chronic severe pain or disabling illness?

- No
 Yes

6. Past suicidal behavior in family or associate?

- No
 Yes

RECENT EVENTS (within the past 3 months)

1. Recent significant loss?

- No
 Yes

2. Recent psychiatric admission or discharge?

- No
 Yes

3. Recent first diagnosis of any psychiatric disorder?

- No
 Yes

4. Recent worsening of depressive symptoms or increase in alcohol abuse?

- No
 Yes

CURRENT STATUS (within the last week)

1. Current preoccupation and plans for suicide?

- No
 Yes

2. Current psychomotor agitation or marked anxiety?

- No
 Yes

3. Current prominent feelings of hopelessness?

- No
- Yes

4. Currently living alone?

- No
- Yes

Clinician's Rating of the level of concern about potential suicidal behavior:

What is your level of concern about potential suicidal behavior for this patient?

- Lowest concern (no prior or current concern about suicidal behavior)
- Some concern (prior history of suicidal ideation or behavior, but preventing suicidal behavior is not a focus of the current clinical management)
- Moderate concern (preventing suicidal behavior is part of current clinical management of the patient)
- High concern (preventing suicidal behavior is one of the main goals in the current management of the patient)
- Imminent concern (preventing suicidal behavior is the most important goal in the current clinical management of the patient)

WHODAS Clinician-Rated Disability Assessment

Please answer the questions below based on the overall clinical presentation of the patient and your best clinical judgment after a review of all the information you have obtained during this assessment.						
How do you rate your patient's overall health in the past 30 days?	Very good	Good	Moderate	Bad	Very bad	
In the last THIRTY (30) DAYS, how much difficulty did your patient have in:						
1.	Understanding and communicating? That is, - Concentrating or remembering – Finding solutions to problems – Learning something new – Generally understanding and communicating with people	None	Mild	Moderate	Severe	Extreme/ Cannot Do
2.	Getting around? That is, - Standing for long periods – Standing up from sitting down – Moving around inside the home – Getting out of the home – Difficulty with walking a long distance such as a half mile (or kilometer)	None	Mild	Moderate	Severe	Extreme/ Cannot Do
3.	Self Care? That is, - Washing his/her whole body – Getting dressed – Eating – Staying alone for a few days	None	Mild	Moderate	Severe	Extreme/ Cannot Do
4.	Getting along with people? That is, - Dealing with people who are strangers – Maintaining a friendship – Getting along with people who are close – Controlling feelings	None	Mild	Moderate	Severe	Extreme/ Cannot Do
5.	Household activities or work or school activities? That is, - Getting these activities done – Doing these activities well – Doing them as quickly as needed	None	Mild	Moderate	Severe	Extreme/ Cannot Do
6.	Participation in society? That is, - The world and other people creating problems – Discrimination – Problems in living with dignity – Problems in joining in community activities	None	Mild	Moderate	Severe	Extreme/ Cannot Do
Total Score for the 6-item WHODAS:						
Interference with Life: Overall, how much did all of these difficulties interfere with your patient's life?						
	Not at all	Mildly	Moderately	Severely	Extremely	
Form Status:		Incomplete	Unverified	Complete		

Clinician-rated
Diagnostic-Specific Severity Measures

1. Any disorder.....Clinician-rated Overall Severity Disorder
2. Discrepancy.....Clinician/Patient Severity Discrepancy
3. Anxiety.....Behavioral Severity Dimensions: Anxiety
4. Psychosis.....Diagnostic-Specific Severity Measure for Psychosis

Clinician-rated Overall Severity

of fill in disorder

Clinician-rated Overall Severity of _____

(Please fill in disorder)

Based on your overall clinical evaluation including your review of the patient's symptom presentation as indicated in the patient-rated cross-cutting dimensional measures and the patient-rated severity measure, please rate the overall SEVERITY of the patient's _____ disorder.

(Note: The overall severity of the patient's condition is based on your clinical judgment)

Please check a box below:

- (0) None
- (1) Slight
- (2) Mild
- (3) Moderate
- (4) Severe

Discrepancy:

If there is discrepancy between your rating of the severity of the patient's condition and the patient's perception of the severity of his/her condition, please provide a brief explanation for your decision:

Behavioral Severity Dimensions: Anxiety

Instructions: Clinicians, please complete if the patient meets criteria for any mood disorder.

Anxious Symptoms:

- a. describes (irrational) worries
- b. feeling uneasy
- c. feeling nervous
- d. motor tension
- e. feels something awful may happen

Anxious Distress - defined as 3 or more of the above anxious symptoms.

Clinical Anxiety Scale

- Not anxious
- Mildly anxious
- Moderate anxiety – 2 symptoms
- Severely anxious 3–5 symptoms
- Severely anxious with motor agitation

Diagnostic-Severity Measure for Psychosis

Instructions: Clinicians, please complete if the patient reaches diagnostic criteria for any psychotic disorders. The dimensional domains are assessed on a 0-4 scale on a cross-sectional basis, with severity assessment based on **the past month**.

	Hallucinations	Delusions	Disorganization	Abnormal Psychomotor Behavior	Restricted Emotional Expression	Avolition	Impaired Cognition	Depression	Mania
0	Not Present	Not Present	Not Present	Not Present	Not Present	Not Present	Not Present	Not Present	Not Present
1	Equivocal (severity or duration not sufficient to be considered psychosis)	Equivocal (severity or duration not sufficient to be considered psychosis)	Equivocal (severity or duration not sufficient to be considered disorganization)	Equivocal (severity or duration not sufficient to be considered abnormal psychomotor behavior)	Equivocal decrease in facial expressivity, prosody, or gestures	Equivocal decrease in self-initiated behavior	Equivocal (cognitive function not clearly outside the range expected for age and SES, i.e., within 1 SD of mean)	Equivocal (some depressed mood, but insufficient symptoms, duration or severity to meet diagnostic criteria)	Equivocal (some inflated or irritable mood, but insufficient symptoms, duration, or severity to meet diagnostic criteria)
2	Present, but mild (little pressure to act upon voices, not very bothered by voices)	Present, but mild (delusions are not bizarre, or little pressure to act upon delusional beliefs, not very bothered by beliefs)	Present, but mild (some difficulty following speech and/or occasional bizarre behavior)	Present, but mild (occasional abnormal motor behavior)	Present, but mild decrease in facial expressivity, prosody, or gestures	Present, but mild in self-initiated behavior	Present, but mild (some reduction in cognitive function below expected for age and SES, b/w 1 and 2 SD from mean)	Present, but mild (meets criteria for Major Depression, with minimum number of symptoms, duration, and severity)	Present, but mild (meets criteria for Mania with minimum number of symptoms, duration, and severity)
3	Present and moderate (some pressure to respond to voices, or is somewhat bothered by voices)	Present and moderate (some pressure to act upon beliefs, or is somewhat bothered by beliefs)	Present and moderate (speech often difficult to follow and/or frequent bizarre behavior)	Present and moderate (frequent abnormal motor behavior)	Present and moderate decrease in facial expressivity, prosody, or gestures	Present and moderate in self-initiated behavior	Present and moderate (clear reduction in cognitive function below expected for age and SES, b/w 2 and 3 SD from mean)	Present and moderate (meets criteria for Major Depression with somewhat more than the minimum number of symptoms, duration, and/or severity)	Present and moderate (meets criteria for Mania with somewhat more than the minimum number of symptoms duration, and/or severity)
4	Present and severe (severe pressure to respond to voices, or is very bothered by voices)	Present and severe (severe pressure to act upon beliefs, or is very bothered by beliefs)	Present and severe (speech almost impossible to follow and/or behavior almost always bizarre)	Present and severe (abnormal motor behavior almost constant)	Present and severe decrease in facial expressivity, prosody, or gestures	Present and severe in self-initiated behavior	Present and severe (severe reduction in cognitive function below expected for age and SES, > 3SD from mean)	Present and severe (meets criteria for Major Depression with many more than the minimum number of symptoms and/or severity)	Present and severe (meets criteria for Mania with many more than the minimum number of symptoms and/or severity)